

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

If we turn to the nucleus, we find similarly that the chromomeres, the chromosomes, and the leaders are also ephemeral and secondary structures which form and disperse, the chromioles alone being the permanent individualities of the chromosomic structures. The nucleus then contains the following permanent granula: linosomes, the chromoplasmic granula, the chromioles, and the granula composing the chromoplasts. The permanent structures of the cell are the centrioles, the chromioles, and the chromoplasts. As regards the primary parts of these last-mentioned structures we are yet in doubt, but there is every reason to believe that these structures are of a highly complicated nature.

Similarly the fibrillar and alveolar structures of the protoplasm are only secondary, ephemeral, and temporary. With proper optical means we see that the alveole, as well as the reticulum, is built up of granules. These granules adhere to each other by means of minute projections or arms, for which I have proposed the name of Linopodia. The ultimate visible structure of the protoplasm is thus a granule, capable of projecting and retracting Linopodia.

For a fuller explanation and demonstration of these facts I must refer to the larger paper now in the hands of the publisher of the *Journal of Morphology*.

BIOLOGICAL LABORATORY,

CALIFORNIA ACADEMY OF SCIENCES,

SAN FRANCISCO, CALIFORNIA.

ERRATUM.

In No. 1, p. 1, 13th line, last word, read ganglion in place of "gland."